



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Appropriate selection of convalescent plasma donors for COVID-19

We read with considerable interest the Comment from Long Chen and colleagues¹ about the potential use of convalescent plasma for the treatment of COVID-19. Chen and colleagues mention the earlier pragmatic WHO recommendation for the use of convalescent plasma as therapy in Ebola virus disease.² The absence of a clinically relevant therapeutic benefit in patients with Ebola virus infection described by Griensven and colleagues,³ and more recently the finding of no therapeutic benefit in a small trial in patients with COVID-19 in Zhengzhou, China,⁴ will be used to question the usefulness of convalescent plasma in COVID-19. In the Guinea-Bissau Ebola study,³ no attempt was made to select donors for the potency of their neutralising antibody. In the COVID-19 study,⁴ seropositive donors were recruited only after IgM antibody to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was no

longer detected, and no attempt to quantify the antibody response was reported. We previously described the levels of detectable antibody and the inferred level of neutralising antibody in convalescent plasma donors for patients with Ebola virus disease in Sierra Leone,⁵ showing 100-fold differences in the level of neutralising antibody. We described a strategy for selecting donors with the highest levels of neutralising antibody, which was not undertaken in donors in the Guinea-Bissau or Zhengzhou studies. For planned interventions in the treatment of patients with COVID-19 severe disease, we strongly recommend selection and qualification only of donors who carry the highest levels of detectable neutralising antibody to SARS-CoV-2. In this respect, we have data which indicate that quantification of specific antibody to the receptor-binding domain will indicate levels of neutralising antibody (unpublished). Commercial assays based on the receptor-binding domain alone, although not intended for the purpose of identifying suitable convalescent plasma donors, will probably be able to serve this need.

MGS reports grants from DHSC National Institute for Health Research, Medical Research Council UK, and Health Protection Research Unit in Emerging and Zoonotic Infections at the University of Liverpool during the conduct of this work and being a minority owner of Integrum Scientific outside the submitted work. RST declares no competing interests.

*Richard S Tedder, Malcolm G Semple
r.tedder@imperial.ac.uk

Department of Infectious Disease, Imperial College London, London W21PG, UK (RST); and Health Protection Research Unit in Emerging and Zoonotic Infection, University of Liverpool, Liverpool, UK (MGS)

- 1 Chen L, Xiong J, Bao L, Shi Y. Convalescent plasma as a potential therapy for COVID-19. *Lancet Infect Dis* 2020; **20**: 398–400.
- 2 WHO. Use of convalescent whole blood or plasma collected from patients recovered from Ebola virus disease for transfusion, as an empirical treatment during outbreaks. 2014. <https://www.who.int/csr/resources/publications/ebola/convalescent-treatment/en> (accessed June 3, 2020).
- 3 van Griensven J, Edwards T, de Lamballerie X, et al. Evaluation of convalescent plasma for Ebola virus disease in Guinea. *N Engl J Med* 2016; **374**: 33–42.
- 4 Zeng Q-L, Yu Z-J, Gou J-J, et al. Effect of convalescent plasma therapy on viral shedding and survival in COVID-19 patients. *J Infect Dis* 2020; published online April 29. DOI:10.1093/infdis/jiaa228.
- 5 Tedder RS, Samuel D, Dicks S, et al. Detection, characterization, and enrolment of donors of Ebola convalescent plasma in Sierra Leone. *Transfusion* 2018; **58**: 1289–98.



Lancet Infect Dis 2020

Published Online

June 15, 2020

[https://doi.org/10.1016/S1473-3099\(20\)30470-9](https://doi.org/10.1016/S1473-3099(20)30470-9)